

Medihaler-ISO®


Medihaler-EPI®

Isoproterenol sulfate, 2.0 mg. per cc., suspended in inert, nontoxic aerosol vehicle. Contains no alcohol. Each measured dose contains 0.06 mg. isoproterenol.

Epinephrine bitartrate, 7.0 mg. per cc., suspended in inert, nontoxic aerosol vehicle. Contains no alcohol. Each measured dose contains 0.15 mg. epinephrine.

NOTABLY WELL TOLERATED AND EFFECTIVE FOR CHILDREN, TOO







In hypoprothrombinemia

KONAKION

Rapid action

Wide margin of safety

Versatility of administration

Compatibility

Low dosage forms

rate of absorption faster than menadione or derivatives...more potent and lasting effects. substantially safer than vitamin-K analogues-no

substantially safer than vitamin-K analogues—no kernicterus reported.

capsules for oral use... fine aqueous dispersion for parenteral administration.

unlike vitamin-K analogues or similar products, the parenteral form of Konakion is a fine aqueous dispersion compatible with most I.V. vehicles.

no excess, no waste-packaged for economical one-time use.

Prophylactically and therapeutically, Konakion is indicated in obstetrics to prevent or control neonatal hemorrhage, to minimize excessive bleeding in surgery, to offset anticoagulant overdosage, and whenever vitamin-K utilization is impaired.

KONAKION®-brand of vitamin K1

Capsules-5 mg; Ampuls-1 mg/0.5 cc



ROCHE LABORATORIES • Division of Hoffmann-La Roche Inc • Nutley 10, New Jersey

The Bulletin of The New York Academy of Medicine, Vol. 35, No. 6, June, 1959. Published Monthly by The New York Academy of Medicine. 2 East 103 Street, New York 29, N. Y. Entered as second-class matter February 3, 1928, at the Post Office at New York, N. Y., under the act of August 24. 1912. Postage paid at New York, N. Y. Annual subscription United States and Canada \$8.00. All other countries \$9.00. Single copies 90c.



Butazolidin

tablets · alka capsules

potent · nonhormonal · anti-inflammatory agent

BUTAZOLIDIN tablets or the Alka capsules are equally effective but individually adaptable in a wide range of arthritic disorders.

Recent clinical reports continue to justify the selection of Butazolidin for rapid relief of pain, increased mobility, and early resolution of inflammation.

Gouty Arthritis: "...95 per cent of patients experienced a satisfactory response..."

Rheumatoid Arthritis: In "A total of 215 cases...over half, 50.7 per cent showed at least major improvement, with 21.8 per cent showing minor improvement...."² Osteoarthritis: 301 cases showed "...a total of 44.5 per cent with complete remission or major improvement. Of the remainder, 28.2 per cent showed minor improvement...."² Spondylitis: All patients "...experienced initial major improvement that was maintained throughout the period of medication."³ Painful Shoulder Syndrome: Response of 70 patients with various forms showed "...8.6 per cent complete remissions, 47.1 per cent major improvement, 20.0 per cent minor improvement..."²

References: 1. Graham, W.: Canad. M. A. J. 79:634 (Oct. 15) 1958. 2. Robins, H. M.; Lockie, L. M.; Norcross, B.; Latona, S., and Riordan, D. J.: Am. Pract. Digest Treat. 8:1758, 1957. 3. Kuzell, W. C.; Schaffarzick, R. W.; Naugler, W. E., and Champlin, B. M.: New England J. Med. 256:388, 1957.

Availability BUTAZOLIDIN® (phenylbutazone egicy): Red coated tablets of 100 mg. BUTAZOLIDIN® Alka: Capsules containing BUTAZOLIDIN® (phenylbutazone egicy), 100 mg.; dried aluminum hydroxide gel, 100 mg.; magnesium trisilicate 150 mg.; homatropine methylbromide, 1.25 mg.



studies in 706 patients show that

DESTIN



promotes "early, clean

and healthy healing"1

in || traumatic and infectious wounds

- burns (first, second, third degree)
- abdominal fistulae and wounds
- pressure sores and ulcers
- pilonidal cysts and sinuses
- ano-rectal wounds chest wounds samples and new reprint on request.

DESITIN CHEMICAL COMPANY

812 Branch Ave., Providence 4, R. I.

1. Grayzel, H. G., and Schapiro, S.: Western J. Surgery, Obstet. & Gyn., Oct. 1956.

S.: J.A.M.A. 157:1594, April 30, *1. Borrus, J. C. J.A.M.A. 157:159 il 30. B. and coutti, C.: Lav. neuro-th. 17:7 March 1956. 6. Berglund, ocutti, C.: Lav. neuro-1955. 3. Lemere, F .: thwest Med 98, Oct. 4. Bene psichiat. 18:693. 1 5. Sellin .: J. Clin Exper. Psy ska läk. <u>5</u> 3362, Dec. 1956. 7 el, H. A., Wood, J. A. M., Blumenthal, B. lordh, P.: C., Elkins, H., Hiler, .: Rass. med. 34:233, . 67:780, May 9, 1957. ollister and Dixon, H. H.: York Acad Ann. New Yo Acad. Sc. 67:789, May 9, 7. 9. G E. G. and St. Pierr Niklund, P exis 20:27, J. Pediat. B 1957 July-Aug 57. 10 1, R. et al. 7, Aug 12. Tucker, W. I.: South . J. <u>50</u>:1111, Sept. 193 3. Rolandi thesist 6 ecker, W.: Klin. Mon Nov.-Dec. 1957, 14. 1. Augenh 2):224, ostet'. 3(6 ebianco G. and Ce 9:1914, 1958. 17. G. and Ceroni, T.: Oto-r 26(2):143, 1958. 16. o, G.: 1958. ing. ital. Minery tuso, R. and ti, F.: Ac eurol. 3(1):36, J 1953. 49, Feb. 105 astur, R. : J. Indian M. Prof 13) Feb. 18. 1 Gazz, med. ital. .: Lyon méd. <u>19</u> 1958. f, W.: Therap. 97:66, Fe 21. Be March 2. 1958. iglia, G.: Mine 10:218, M 15, 1958. lander, H. S.: Cardiol. 75:101,1 25. Sprauer, V. 1958. 24. McClen Arch. Ped Internat. 1:395. GP Clinics 171: 1958. 26. pinson, R. Robinson, H. M., r.: South. ed. <u>58</u>:1285, April 15, 1958. M. J. 51.5 9, April 1958. 27. S. and Pe er, L.: Ne 18 1958. 29. Reboul, E., Reboul, e: Ann. méd. ychol. <u>116</u> 28. Bouquerel, J., Naviau and M. and Dorgeuille, C.: Maroc 37:784, July 1958. 30. B Lyon med. 200:885, Nov. 1958. Med. Klin. 53:2113, Dec. 1958. 31. Lamphier, T. A.: Maryland M 7:627, Nov. 1958. 32. Led

70-95%

of nervous, tense patients recovered or improved

For your patients, Miltown promptly checks emotional and muscular tension. Thus, you will make it easier for them to lead a normal family life and to carry on their usual work.

For you, the choice of Miltown as the tranquilizer means the comfortable assurance that it will relieve nervousness and tension without impairing your patient's mental efficiency, motor control, normal behavior or autonomic balance.





the pattern of

GLUCOSAMINE-POTENTIATED TETRACYCLINE

therapy

COSA-TETRACYN*

125 mg., 250 mg.


oral suspension orange flavored, 2 oz. bottle, 125 mg. per teaspoonful (5 cc.)

pediatric drops orange flavored, 10 cc. bottle (with calibrated dropper), 5 mg. per drop (100 mg. per cc.)

Plizer Science for the world's well-being PFIZER LABORATORIES

Division, Chas. Pfizer & Co., Inc. Brooklyn 6, N. Y.

*Trademark for glucosamine-potentiated tetra cycline



specify Lactum a well tolerated formula ... widely and satisfactorily used

Normal Stool Frequency with Lactum¹

Lactum was fed to 57 normal infants for 2 to 10 months. The investigators observed: "In no case was there any history of persistent diarrhea. In forty-eight infants, the usual number of stools was one to two daily; in only nine were there three, or occasionally four a day."

Low Incidence of Loose Stools in Hospital Study2

In a clinical study of 180 sick and 10 well infants fed Lactum, the investigator reported:
"There was no diarrhes in any of the infants "There was no diarrhea in any of the infants except in those patients whose diagnosis on admission was infectious diarrhea... we have been convinced that inclusion of adequate added carbohydrate in ... [Lactum] during stress periods, such as diarrhea, is definitely advantageous."

Age	No. Cases	Loose Stools†
Birth	14	0
1 mo.	36	0
2 mo.	27	1
3 mo.	12	1
4 mo.	10	1
5 mo.	14	0
6 mo.	8	0
7 mo.	19	0
8 mo.	11	0
9 mo.	10	0
10 mo.	8	0
11 mo.	11	0
12 mo.	10	0
	190	3 (1.6

†Exclusive of infants admitted with infectious diarrhea.

1. Frost, L. H., and Jackson, R.L.: J. Pediat. 39: 585-592 (Nov.) 1951.

2. Henrickson, W. E.: GP 8:51-56 (Oct.) 1953.

to avoid baby's discomfort and mother's anxiety...specify

actum liquid · "instant" powder Modified milk formula, Mead Johnson





THUS SQUIBB OFFERS YOU GREATER LATITUDE IN SOLVING THE PROBLEM OF

RTENSIO

WITHOUT FEAR OF SIGNIFICANT POTASSIUM DEPLETION 1-3

Rautrax combines Raudixin with flumethiazide - the new, safe nonmercurial diuretic — for control of all degrees of hypertension. Clinicians report it safely and rapidly eliminates excess extracellular sodium and water without potassium depletion.¹⁻³ Potassium loss is less than with any other nonmercurial diuretic.1 Moreover, the inclusion of supplemental potassium chloride in Rautrax provides added protection against potassium and chloride depletion in the long-term management of hypertension.

Through this dependable diuretic action of flumethiazide, the clinical and subclinical edema - so often associated with cardiovascular disease — is rapidly brought under control.2-5 And once Rautrax has brought the fluid balance within normal limits, continued administration does not appreciably alter the normal serum electrolyte pattern. Flumethiazide also potentiates the antihypertensive action of Raudixin. By this unique dual action, a lower dosage of each ingredient effectively maintains safe antihypertensive therapy.

Dosage: 2 to 6 tablets daily in divided doses initially; may be adjusted within range of 1 to 6 tablets daily in divided doses. Note: In hypertensive patients already on ganglionic blocking agents, veratrum and/or hydralazine, the addition of Rautrax necessitates an immediate dosage reduction of these agents by at least 50%. A similar reduction is necessary when these agents are added to the Rautrax regimen.

Supply: Capsule-shaped tablets supplying 50 mg, of Raudixin, 400 mg, of flumethiazide, and 400 mg, of potassium chloride, bottles of 100. References: 1. Moyer, J. H., and others: Am. J. Cardiol., 3:113 (Jan.) 1959. • 2. Bodi, T., and others: To be published, Am. J. Cardiol., (April) 1959 • 3. Fuchs, M., and others: Monographs on Therapy, 4:43 (April) 1959. • 4. Montero, A. C.; Rochelle, J. B., III, and Ford, R. V.: To be published. • 5. Rochelle, J. B., III; Montero, A. C., and Ford, R. V.: To be published.

LITERATURE AVAILABLE ON REQUEST.

RAUDIXIN® AND RAUTRAX ARE SQUIBB TRADEMARKS





while she is planning her family,

she needs your help more than ever



the most widely prescribed contraceptive



WHENEVER A DIAPHRAGM IS INDICATED

in epilepsy prerequisite for emotional adjustment: therapy

The most effective form of psychotherapy is to demonstrate to the patient that his seizures can be adequately controlled by the use of anticonvulsant medication."*

requisite for therapy:

THE PARKE-DAVIS FAMILY OF ANTICONVULSANTS

effective anticonvulsants for most clinical needs

for control of grand mal and psychomotor seizures

Diantin [®] Sodium (diphenylhydantoin sodium, Parke-Davis) is supplied in several forms including Kapseals of 0.03 Gm. and of 0.1 Gm. in bottles of 100 and 1,000.

Phe antin [®]Kapseals (Dilantin 100 mg., phenobarbital 30 mg., desoxyephedrine hydrochloride 2.5 mg.), bottles of 100.

for the petit mal triad Kapseals (phensuximide, Parke-Davis) 0.5 Gm., bottles of 100 and 1,000. Suspension, 250 mg. per 4 cc., 16-ounce bottles.

Ceontin[®] Kapseals (methsuximide, Parke-Davis) 0.3 Gm., bottles of 100.

Literature supplying details on dosage and administration available on request. *Carter, S.: M. Clin. North America: 315 (March) 1953.






TRICHOMONAS MONILIA BACTERIA

welcome clinical advance...

effective medication in an appealing form



Soft and pliant as a tampon, the Milibis vaginal suppository offers proved therapeutic action* in a vehicle giving unusual clinical advantages to both patients and physician.

COVERS CERVIX AND VAGINAL WALL—The pliant Milibis suppository disintegrates readily and molds itself to the cervix as well as the columns and rugae of the vaginal vault.

SHORT DOSAGE SCHEDULE—The short course of treatment with Milibis—only 10 suppositories in most cases—together with the clean, odorless, non-staining qualities eliminates psychic barriers which often interrupt longer treatments before complete cure.

MILIBIS® Vaginal Suppositories

Now supplied with plastic applicator

- SANITARY
 - INSURES CORRECT SUPPOSITORY PLACEMENT

Winthrop LABORATORIES
Now York 18, N.Y.

*97 per cent effective in a study of 564 cases; 94 per cent effective in a series of 510 cases.

Milibis (brand of glycobiarsol), trademark reg. U.S. Pat. Off.

SUPPLIED: BOXES OF 10 with applicator.



both blood picture and patient respond to $\textbf{TRINSICON}^\circ$

Investigators^{1,2} have determined that low serum iron may be accompanied by insidious vitamin B_{12} deficiencies which result from subnutrition, increased demand, or lack of intrinsic factor. Coexisting vitamin C deficiencies also have been found.³


These studies suggest that an anemia may be multiple in nature—that optimum results would be derived from a combination of therapeutic agents.

Trinsicon offers therapeutic quantities of all known hematinic factors. Prescribe two Pulvules® daily to provide assured response in all treatable anemias.

Trinsicon® (hematinic concentrate with intrinsic factor, Lilly)

1. A. M. A. Arch. Int. Med., 99:346, 1957. 2. Am. J. Obst. & Gynec., 70:1309, 1955. 3. Lancet, 1:448, 1957.

ELI LILLY AND COMPANY . INDIANAPOLIS 6, INDIANA, U.S.A.





Underweight Children Gain and Retain Weight with Nilevar*

One of the most convincing evidences of the anabolic activity of Nilevar, brand of norethandrolone, has been its ability to improve appetite and increase weight in poorly nourished, underweight children.

A highly important feature of the weight gain thus produced is that it is not ordinarily manifested by deposition of fat but as muscle tissue resulting from the protein anabolism induced by Nileyar.

Anorexia and "Weight Lag" Study—Brown, Libo and Nussbaum have reported* consistent and definite increases in rate of weight gain in eighty-six patients, ranging in age from 7 weeks to 15½ years. This beneficial action of Nilevar was observed in the patients with organic and traumatic disorders as well as those whose only complaints were poor appetite and/or persistent failure to gain weight.

In this study, the weight gained was not lost

after discontinuance of Nilevar therapy although many patients did not continue the sharp gains effected by the drug.

The authors are of the opinion that Nilevar is a highly useful anabolic agent for influencing weight gain in underweight children.

When Nilevar is administered to children a dose of 0.25 mg. per pound of body weight is recommended and continuous dosage for more than three months is not recommended.

Nilevar is supplied as tablets of 10 mg., drops of 0.25 mg. per drop and ampuls of 25 mg. in 1 cc. of sesame oil. Further dosage information in Searle Reference Manual No. 4.

G. D. Searle & Co., Chicago 80, Illinois. Research in the Service of Medicine.

^{*}Brown, S. S.; Libo, H. W., and Nussbaum, A. H.: Norethandrolone in the Successful Management of Anorexia and "Weight Lag" in Children, Scientific Exhibit presented at the Annual Meeting of the American Academy of Pediatrics, Chicago, Oct. 20-23, 1958.

for total of itc infecte skin lesions



Cleared in 5 days

Kenalog, Spectrocin and Mycostatin in Plastibase


antipruritic/anti-inflammatory/antibacterial/antifungal

Mycolog Ointment - containing the new superior topical corticoid Kenalog - reduces inflammation,3.4 relieves itching,1.2 and combats or prevents bacterial, monilial and mixed infections. 5-7 It is extremely well tolerated, and assures a rapid, decisive clinical response for most infected dermatoses.

"Thirty-one of 38 patients . . . obtained excellent or good control of dermatological lesions . . . [Mycolog] was highly effective, particularly in the management of mixed infections. Several recalcitrant eruptions which had not responded to previous therapy were remarkably responsive to the daily application of this preparation over periods of 2 to 3 weeks."5

For total management of itching, inflamed, infected skin lesions, Mycolog contains triamcinolone acetonide, an outstanding new topical corticoid for prompt, effective relief of itching, burning and inflammation1-4 - neomycin and gramicidin for powerful antibacterial action7 - and nystatin for treating or preventing Candida (Monilia) albicans infections.8,9

Application: Apply 2 to 3 times daily. Supply: 5 Gm. and 15 Gm. tubes. Each gram supplies 1.0 mg. (0.1%) triamacetonide, 2.5 mg. neomycin base, 0.25 mg. gramicidin, and 100,000 units nystatin in Plastibase References: 1. Shelmire, J.B., Jr.: Monographs on Therapy 3:164 (Nov.) 1958. • 2. Nix, T.E., Jr., and Derbes, V.J.: Monographs on Therapy 3:123 (Nov.) 1958. • 3. Robinson, R.C.V.: Bull. School of Med., U. Maryland 43:54 (July) 1958. • 4. Sternberg, T.H.: Newcomer, V.D., and Reisner, R.M.: Monographs on Therapy 3:115 (Nov.) 1958. • 5. Clark, R.F., and Hallett, J.J.: Monographs on Therapy, 3:153 (Nov.) 1958. • 6. Smith J.G., Jr.; Zawisza, R.J., and Blank, H.: Monographs on Therapy, 3:111 (Nov.) 1958. • 7. Monographs on Therapy, 3:137 (Nov.) 1958. • 8. Howell, C.M., Jr.: North Carolina M.J. 19:449 (Oct.) 1958. • 9. Bereston, E.S.: South, M.J. 50:547 (April) 1957. And whatever the topical corticoid need, a suitable Squibb formulation is available—Kenalog-S Lotion—7½ cc. plastic squeeze bottles. Each cc. supplies 1.0 mg. (0.1%) triamcinolone acetonide, 2.5 mg. neomycin base and 0.25 mg, gramicidin. Kenalog Cream, 0.1%—5 Gm. and 15 Gm. tubes. Kenalog Lotion, 0.1%—15 cc. plastic squeeze bottles. Kenalog Cintment, 0.1%—5 Gm. and 15 Gm. tubes.



Cleared in 20 days



SQUIBB

Squibb Quality — the Priceless Ingredient

"SPECTROCIN"B, "MYCOSTATIN"B, "PLASTIBASE"B, "MYCOLOG"



the completely new, 2-speed

SANBORN Model 100 VISO-CARDIETTE

Here is an electrocardiograph in which no detail has been overlooked to give you diagnostically accurate information ... the greatest possible operating convenience... and modern, functional attractiveness. With thirty-five years of experience, this is the finest electrocardiograph Sanborn Company has ever produced. Priced at eight hundred fifty dollars, delivered continental U.S.A.

SANBORN COMPANY

MEDICAL DIVISION • 175 Wyman St., Waltham 54, Mass.

NEW YORK Branch Office 1841 Broadway, Circle 7-5794 and 7-5795

There is no Penicillin in Walker-Gordon Certified Milk

Exceptional technical control measures assure the physician that Walker-Gordon Certified Milk is free of all penicillin and other antibiotic residue.

This control has recently been incorporated by The American Association of Medical Milk Commissions in its "Methods and Standards for the Production of Certified Milk."

Patients cannot develop any unfavorable antibiotic reaction as a result of drinking Walker-Gordon Certified Milk.



For more detailed information write or phone...

Walker-Gordon Certified Milks

Farm: Plainsboro, N.J. SWinburne 9-1234

New York Office: WAlker 5-7300

Philadelphia Office: LOcust 7-2665
Certifled by Medical Milk Commissions of

 CERTIFIED
 New York, Kings, Hudson, and Philadelphia Counties

 Walker Gordon Certified Milks ★ Raw ★ Past. ★ Homo. ★ Skimmed ★ Lo-Sodium ★ Acidophilus

For acidosis due to anesthesia — edema



KALAK

Counter-Acts ANTI-BIOTIC REACTIONS

. . . . KALAK is a nonlaxative, alkaline diuretic buffer — side reactions from aureomycin — terramycin — sulfas — penicillin are reduced through the use of KALAK — KALAK

contains only those salts NORMALLY present in plasma. . . . IT IS BASIC!

KALAK WATER CO. of NEW YORK, Inc.

207 EAST 37TH STREET New York 16, N. Y.

For acidosis due to nausea — in nephritis

BULLETIN

THE NEW YORK ACADEMY OF MEDICINE


 \Diamond

The Bulletin contains scientific papers presented at the Academy, including . . .

Post-Graduate Week Lectures
Stated Meeting Addresses of the Academy
Academy Section Papers
Section on Microbiology—Abstracts
Monthly Panel Meetings—Transcripts
New York Pathological Society—Abstracts
and Other Original Contributions
Published Monthly

Annual subscription price, \$8.00 in United States and Canada. All other countries \$9.00

for the peak of analgesic efficiency



DILAUDID

Dosage Forms of Dilaudid hydrochloride:

Ampules: 1cc., 2 mg. and 3 mg. each. Hypodermic Tablets: 2, 3 and 4 mg. each.

Oral Tablets: 2.7 mg. each.

Multiple Dose Vial: 10 cc., 2 mg. Dilaudid sulfate per cc.

*Subject to Federal narcotic regulations Dilaudid®, E. Bilhuber, Inc.

KNOLL PHARMACEUTICAL COMPANY

ORANGE

Improved

THI-LI-FER® FORTE

More than just a hematinic . . . a valuable adjunct in the treatment of subnutritional conditions.

FOR INTRAMUSCULAR USE

FORMULA: A specially stabilized aqueous solution of Cyanocobalamin (Vitamin Bız), highly purified Liver Extract, Thiamine Hydrochloride (Vitamin B₁₎ and Iron.

SUPPLIED: 10 cc. multiple dose vial. No. 383. Also available as THI-LI-FER SYRUP and THI-LI-FER CAPSULES.

C. F. KIRK COMPANY

Pharmaceutical and Biological Laboratories

521 West 23 Street, New York 11

For complete catalog and samples write Dept. 6B

THIASTIGMINE

EFFECTIVE THERAPY FOR PROMPT RELIEF IN THE TREATMENT OF

- NEURITIS
- NEURALGIAS
- HERPES ZOSTER ●

FOR INTRAMUSCULAR USE

FORMULA: Each cc. contains a stabilized aqueous solution of Thiamine Hydrochloride (B1), Neostigmine Methylsulfate, and Atropine Sulfate.

SUPPLIED: 10 cc. multiple dose vial, No. 339.

C. F. KIRK COMPANY

Pharmaceutical and Biological Laboratories

521 West 23 Street, New York 11

For complete catalog and samples write Dept. 6B

Herald Square, N. Y.

Roosevelt Field, Long Island

Parkchester, Bronx



Jamaica, Queens

White Plains, N.-Y.

Flatbush, Brooklyn

MACY'S PRESCRIPTION DEPARTMENT

Macy's respects professional ethics. We carefully avoid discussing your prescription, either its ingredients or its use, with your patient. You can be assured that all your prescriptions are accurately compounded exactly as you write them. And all our preparations, made from the finest ingredients obtainable, are carefully checked by our supervising pharmacists.

No wonder so many physicians prefer Macy's Prescription Department.

THE NEW YORK ACADEMY OF MEDICINE

Third Annual

POSTGRADUATE WEEK 1959

October 5 — 9, inclusive

RESEARCH CONTRIBUTIONS TO CLINICAL PRACTICE

All inquiries should be addressed to: Secretary, Postgraduate Week

THE NEW YORK ACADEMY OF MEDICINE 2 East 103 Street, New York 29, N. Y.

CONTACT service to physicians!

THEO. E. OBRIG. INC.

BRIG, INC.

Is for more than a quarter century
es only upon prescription and
it.
I lenses including Lacrilens®
can be worn for long periods
inded for all sports

N. Y. • PLaza 8-0973 specialists in fitting contact lenses for more than a quarter century Theo. E. Obrig, Inc. fits contact lenses only upon prescription and

under the supervision of a physician.


Expert in fitting all types of contact lenses including Lacrilens® and latest forms of cornea lenses.

Lacrilens® and corneal type lenses can be worn for long periods of time in aphakia.

Lacrilens® is particularly recommended for all sports including swimming.

Careful attention to referred cases.

Theo. E. Obrig, Inc. • 75 East 55th Street, New York 22, N. Y. • PLaza 8-0973 $\overline{\chi}_{ ext{event}}$ when the present of the prese



SIGN OF GOOD TASTE


pyelonephritis

"the most important concept is that it is a tubular disease"1

FURADANTIN

brand of nitrofurantoin

a most important characteristic: effective at the tubular level



In addition to simple glomerular filtration, Furadantin is actively excreted by the cells of the tubules. A significant and singular characteristic of Furadantin, it is but one reason why "the protracted administration of nitrofurantoin [Furadantin] to patients with ineradicable urinary tract infection, particularly chronic pyelonephritis without demonstrable obstruction, may usefully complement the medical management of this difficult problem." Available as Tablets, 50 and 100 mg.; Oral Suspension, 25 mg. per 5 cc. tsp.

References: 1. Smith, I. M., and Lenyo, L.: Am. Practitioner 9:78, 1958. 2. Jawetz, E., et al.: A.M.A. Arch. Int. M. 100:549, 1957.

NITROFURANS—a unique class of antimicrobials—neither antibiotics nor sulfonamides of antimicrobials—neither antibiotics nor sulfonamides

EATON LABORATORIES, NORWICH, NEW YORK

the airway with <u>new</u>
'FEDRAZIL' TABLETS

when colds or allergies congest the respiratory tract

- decongest the entire respiratory tract mucosa
- dilate the bronchi


open

provide potent anti-allergic action

Dosace: Adults and Children over 8 years—1 or 2 tablets, three times daily.

Children 2-8 years—1 tablet daily, or as required.

Each tablet contains: 'Sudafed'® brand Pseudoephedrine Hydrochloride, 30 mg.; 'Perazil'® brand Chlorcyclizine Hydrochloride, 25 mg. Bottles of 100, sugar-coated.



Has There Been a GAP in Your Training?

Are You Capable of Caring for the More than 1 Out of 5 Patients Now Suffering from Easily-Remedied Foot Disabilities?

Do you Know the Competitive Value of Radiotherapy and Surgery for Specific Cancers?

Are You
Confident that
Every Patient
You Treat Is
Completely
Ready to Return
to Normal
Daily Living?

DuVries SURGERY OF THE FOOT

Ingrown nails . . . hammer toes . . . overlapping toes . . . spurs . . . a dozen different kinds of bunions . . . all are very common foot problems, which in spite of their simplicity are very disabling. But, until now complete authoritative information on foot problems like these could not be readily found in the literature. SURGERY OF THE FOOT can help you improve your ability to treat all kinds of foot problems from the minor ingrown nail to the major deformity of a club foot. The only up-to-date book available on this subject, Dr. DuVries' new book gives you both basic principles of foot care and illustrated, step-by-step guidance to treatment of all the surgical foot problems you are likely to encounter. More than 400 drawings, shaded drawings, photographs and roentgenograms add much to the help that this new book can give you.

By HENRI L. DuVRIES, M.D., B.Sc., D.S.C., Clinical Instructor in Surgery, Chicago Medical School; Attending Surgeon, Columbus Hospital, Mother Cabrini Hospital, and Frank Cuneo Hospital, Chicago. Just Published. Approx. 494 pages, 634" x 934", 403 illustrations. Price, \$12.50.

Moss THERAPEUTIC RADIOLOGY

Here for the first time is a book that gives you an up-to-date clinical presentation of modern concepts in radiobiology. It explains current thinking on the behavior of cancer and gives you a practical clinical orientation on the management of specific cancers.

Throughout every chapter of this new book—in every discussion of treatment for all of the more common lesions of nearly every organ and body tissue, you'll find stimulating discussions detailing the opinions of leading radiologists, pathologists and surgeons regarding the values of irradiation and surgery. In addition, you'll have at your fingertips a wealth of practical information on the day-to-day care of your cancer patients.

By WILLIAM T. MOSS, M.D., Director, Department of Therapeutic Radiology, Chicago Wesley Memorial Hospital, Chicago, Ill.; Assistant Professor, Department of Radiology, Northwestern University School of Medicine, Chicago, Ill. Just Published. 403 pages, 634" x 934", 146 figures. Price, \$12.50.

Rusk REHABILITATION MEDICINE

You can be confident that every patient you treat receives this "last step" in total patient care if you will but apply the principles of rehabilitation in your own practice.

of rehabilitation in your own practice.

The new Mosby book, REHABILITATION MEDICINE, can give you an understanding of advanced principles and can show you exactly how to apply them in your work. Written by "Dr. Rehabilitation" himself, Howard A. Rusk, M.D., with the collaboration of 36 of his colleagues in the Department of Physical Medicine and Rehabilitation, New York University Bellevue Medical Center, this 572 page volume is the only comprehensive, up-to-date book which describes modern concepts and practices as they are employed at this world renowned medical center.

By HOWARD A. RUSK, M.D., Professor and Chairman of the Department of Physical Medicine and Rehabilitation, New York University-Bellevue Medical Center, New York City; and 36 Collaborators with the editorial assistance of EUGENE J. TAYLOR, A.M. New. 1958, 572 pages, 634" x 934", 172 figures. Price, \$12.00.

Purchase Any of Three Ways:

At Your Favorite Bookstore; From Our Personal Sales Representative; Or Order on 10 Day Approval From

The C. V. Mosby Company
3207 Washington Boulevard • St. Louis 3, Missouri

